arXiv:1509.04542v2 [math.CA] 7 Jan 2016

Asymptotic zero distribution of Jacobi-Piñeiro and multiple Laguerre polynomials

Thorsten Neuschel, Walter Van Assche
Katholieke Universiteit Leuven, Belgium

January 8, 2016

Abstract

We give the asymptotic distribution of the zeros of Jacobi-Piñeiro polynomials and multiple Laguerre polynomials of the first kind. We use the nearest neighbor recurrence relations for these polynomials and a recent result on the ratio asymptotics of multiple orthogonal polynomials. We show how these asymptotic zero distributions are related to the Fuss-Catalan distribution.

1 Introduction and main results

In this paper we obtain the asymptotic distribution of the zeros of two families of multiple orthogonal polynomials: the Jacobi-Piñeiro polynomials and the multiple Laguerre polynomials of the first kind [9, Ch. 23], [2], [19]. These are two families of multiple orthogonal polynomials for which explicit formulas are known and which are useful for a number of applications. For instance, the zeros of Jacobi-Piñeiro polynomials (and Wronskian-type determinants of Jacobi-Piñeiro polynomials) form the unique solution of certain Bethe Ansatz equations [14] and multiple orthogonal polynomials are also useful for investigating determinantal point processes [10]. Recently the Jacobi-Piñeiro ensemble and the multiple Laguerre ensemble were introduced for random matrix minor processes related to percolation theory [1] which are based on the Jacobi-Piñeiro and multiple Laguerre polynomials of the first kind.

Let $\vec{n} = (n_1, n_2, \ldots, n_r) \in \mathbb{N}^r$ be a multi-index of size $|\vec{n}| = n_1 + n_2 + \cdots + n_r$. The Jacobi-Piñeiro polynomials $P_{\vec{n}}$, with parameters $\vec{\alpha} = (\alpha_1, \ldots, \alpha_r)$ and β, are type II multiple orthogonal polynomials on $[0,1]$ for r Jacobi weights, i.e., $P_{\vec{n}}$ is a monic polynomial of degree $|\vec{n}|$ satisfying

$$\int_0^1 P_{\vec{n}}(x) x^k x^{\alpha_j} (1-x)^\beta \, dx = 0, \qquad k = 0, 1, \ldots, n_j - 1,$$

for $j = 1, 2, \ldots, r$, where $\beta > -1$ and $\alpha_j > -1$ for $1 \leq j \leq r$. They were introduced by Piñeiro for $\beta = 0$ [17]. A multi-index $\vec{n}$ is normal if the monic multiple orthogonal polynomial $P_{\vec{n}}$ of degree $|\vec{n}|$ exists and is unique. All multi-indices for Jacobi-Piñeiro polynomials are normal when $\alpha_i - \alpha_j \notin \mathbb{Z}$ because then the measures form an AT-system

[9, §23.1.2]. The polynomials are given by the Rodrigues formula

$$(-1)^{|\vec{n}|}\prod_{j=1}^{r}(|\vec{n}|+\alpha_j+\beta+1)_{n_j}\ (1-x)^{\beta}P_{\vec{n}}(x) \\ = \prod_{j=1}^{r}\left(x^{-\alpha_j}\frac{d^{n_j}}{dx^{n_j}}x^{n_j+\alpha_j}\right)(1-x)^{|\vec{n}|+\beta}, \quad (1.1)$$

where the product of the differential operators can be taken in any order, since these operators are commuting [9, §23.3.2]. Multiple Laguerre polynomials of the first kind $L_{\vec{n}}$ are given by the Rodrigues formula

$$(-1)^{|\vec{n}|}e^{-x}L_{\vec{n}}(x) = \prod_{j=1}^{r}\left(x^{-\alpha_j}\frac{d^{n_j}}{dx^{n_j}}x^{n_j+\alpha_j}\right)e^{-x}, \quad (1.2)$$

where the product of the differential operators can be taken in any order [9, §23.4.1]. If the parameters $\vec{\alpha} = (\alpha_1, \alpha_2, \ldots, \alpha_r)$ are such that $\alpha_i > -1$ for every i and $\alpha_i - \alpha_j \notin \mathbb{Z}$ $(1 \le i, j \le r)$, then all multi-indices are normal and the polynomials satisfy the following orthogonality properties

$$\int_0^\infty L_{\vec{n}}(x)x^k x^{\alpha_j}e^{-x}\,dx = 0, \qquad k = 0, 1, \ldots, n_j - 1,$$

for $j = 1, 2, \ldots, r$. An explicit expression is given by

$$L_{\vec{n}}(x) = \sum_{k_1=0}^{n_1}\cdots\sum_{k_r=0}^{n_r}(-1)^{|\vec{k}|}\frac{n_1!}{(n_1-k_1)!}\cdots\frac{n_r!}{(n_r-k_r)!} \\ \times\binom{n_r+\alpha_r}{k_r}\binom{n_r+n_{r-1}+\alpha_{r-1}-k_r}{k_{r-1}}\cdots\binom{|\vec{n}|-|\vec{k}|+k_1+\alpha_1}{k_1}x^{|\vec{n}|-|\vec{k}|}. \quad (1.3)$$

We will obtain the asymptotic distribution of the zeros of these multiple orthogonal polynomials by using a result on the asymptotic behavior of the ratio of two neighboring polynomials [21]. This result uses the nearest neighbor recurrence relations for multiple orthogonal polynomials

$$xP_{\vec{n}}(x) = P_{\vec{n}+\vec{e}_k}(x) + b_{\vec{n},k}P_{\vec{n}}(x) + \sum_{j=1}^{r}a_{\vec{n},j}P_{\vec{n}-\vec{e}_j}(x), \qquad 1 \le k \le r,$$

where $\vec{e}_j = (0, \ldots, 0, 1, 0, \ldots, 0)$ with 1 in the jth entry, and some knowledge about the asymptotic behavior for the recurrence coefficients $a_{\vec{n},j}, b_{\vec{n},j}$ $(1 \le j \le r)$. The ratio asymptotic behavior for Jacobi-Piñeiro polynomials will be obtained in Section 2 and for multiple Laguerre polynomials of the first kind in Section 5. The asymptotic distribution of the zeros of Jacobi-Piñeiro polynomials will be obtained in Section 4, where the following result will be proved. We will use the multi-index $\vec{1} = (1, 1, \ldots, 1)$ so that the diagonal index is $(n, n, \ldots, n) = n\vec{1}$.

Theorem 1.1. *Let* $0 < x_{1,rn} < x_{2,rn} < \cdots < x_{rn,rn} < 1$ *be the zeros of the Jacobi-Piñeiro polynomial* $P_{n\vec{1}}$ *with multi-index* $n\vec{1} = (n, n, \ldots, n)$*. Then for every continuous function* f *on* $[0,1]$ *one has*

$$\lim_{n\to\infty} \frac{1}{rn} \sum_{k=1}^{rn} f(x_{k,rn}) = \int_0^1 f(t) v_r(t)\, dt,$$

where the density v_r *on* $[0,1]$ *is given by means of a density* w_r *on* $[0, c_r]$ *as*

$$v_r(x) = c_r w_r(c_r x), \qquad c_r = \frac{(r+1)^{r+1}}{r^r},$$

and with the change of variables

$$\hat{x} = c_r x = \frac{\big(\sin(r+1)\varphi\big)^{r+1}}{\sin\varphi\big(\sin r\varphi\big)^r}, \qquad 0 < \varphi < \frac{\pi}{r+1},$$

the density w_r *is*

$$\begin{aligned} w_r(\hat{x}) &= \frac{r+1}{\pi} \frac{1}{|\hat{x}'(\varphi)|} \\ &= \frac{r+1}{\pi \hat{x}} \frac{\sin\varphi \sin r\varphi \sin(r+1)\varphi}{(r+1)^2 \sin^2 r\varphi - 2r(r+1)\sin(r+1)\varphi \sin r\varphi \cos\varphi + r^2 \sin^2(r+1)\varphi}. \end{aligned} \tag{1.4}$$

The density w_r is in fact the uniform density on $[0, \frac{\pi}{r+1}]$ in the variable φ since

$$\int_0^{c_r} f(\hat{x}) w_r(\hat{x})\, d\hat{x} = \int_0^{\frac{\pi}{r+1}} f(\hat{x}(\varphi)) w_r(\hat{x}(\varphi)) |\hat{x}'(\varphi)|\, d\varphi = \frac{r+1}{\pi} \int_0^{\frac{\pi}{r+1}} f(\hat{x}(\varphi))\, d\varphi.$$

In this sense Theorem 1.1 is the extension to multiple orthogonal polynomials of the equidistribution result for zeros of orthogonal polynomials [18, Thm. 12.7.2] for the case of Jacobi-Piñeiro polynomials. In fact, the same asymptotic distribution of zeros will hold for all multiple orthogonal polynomials for which the nearest neighbor recurrence coefficients behave as in (2.3)–(2.4), provided the zeros of neighboring polynomials interlace. We have plotted the density v_r on $[0,1]$ for $1 \le r \le 5$ in Figure 1.

Observe that

$$\hat{x} = c_r - \binom{r+1}{2} c_r \varphi^2 + \mathcal{O}(\varphi^4), \qquad \varphi \to 0,$$

and

$$\hat{x} = \left(\frac{r+1}{\sin\frac{\pi}{r+1}}\right)^{r+1} \left(\frac{\pi}{r+1} - \varphi\right)^{r+1} + \mathcal{O}\left(\left(\frac{\pi}{r+1} - \varphi\right)^{r+2}\right), \qquad \varphi \to \frac{\pi}{r+1},$$

so that the density v_r behaves as ($\hat{x} = c_r x$)

$$v_r(x) = \mathcal{O}(\varphi^{-1}) = \mathcal{O}\left((1-x)^{-1/2}\right), \qquad x \to 1,$$

and

$$v_r(x) = \mathcal{O}\left(\left(\frac{\pi}{r+1} - \varphi\right)^{-r}\right) = \mathcal{O}\left(x^{-\frac{r}{r+1}}\right), \qquad x \to 0.$$



Figure 1: The asymptotic zero densities v_r for Jacobi-Piñeiro polynomials: $r = 1$ (solid), $r = 2$ (dash), $r = 3$ (dash dot), $r = 4$ (long dash), and $r = 5$ (dots).

Hence the densities v_r have a square root singularity at 1 but a higher order singularity at 0 when $r > 1$, which means that the zeros are more dense near the endpoints 0 and 1, and even more so near 0 than near 1 when $r > 1$. For $r = 1$ the density v_1 is the well-known arcsin density on $[0, 1]$,

$$v_1(x) = \frac{1}{\pi} \frac{1}{\sqrt{x(1-x)}}, \qquad 0 < x < 1,$$

which is the equilibrium measure for $[0, 1]$ in logarithmic potential theory. For $r = 2$ the density can explicitly be written as

$$v_2(x) = \frac{\sqrt{3}}{4\pi} \frac{(1+\sqrt{1-x})^{1/3} + (1-\sqrt{1-x})^{1/3}}{x^{2/3}\sqrt{1-x}}, \qquad x \in (0, 1),$$

and this asymptotic zero distribution was already found in [3, Thm. 2.5]. The moments of w_r are integers given by

$$\int_0^{c_r} x^n w_r(x)\, dx = \frac{r+1}{\pi} \int_0^{\frac{\pi}{r+1}} x(\varphi)^n \, d\varphi = \binom{(r+1)n}{n}, \qquad n \in \mathbb{N} = \{0, 1, 2, \ldots\},$$

which follows from [16, Remark 3.4].

For multiple Laguerre polynomials we need to use a scaling to prevent the zeros from going to infinity. The appropriate scaling is to divide all the zeros of $L_{\vec{n}}$ by $|\vec{n}|$, so that we are in fact investigating the zeros of $L_{n\vec{1}}(rnx)$ for the multi-index $n\vec{1} = (n, n, \ldots, n)$ on the diagonal. In Section 6 we obtain the asymptotic distribution of the scaled zeros, where we prove the following result.

Theorem 1.2. *Let* $0 < x_{1,rn} < x_{2,rn} < \cdots < x_{rn,rn}$ *be the zeros of the multiple Laguerre polynomials* $L_{n\vec{1}}$ *with multi-index* $n\vec{1} = (n, n, \ldots, n)$*. Then for every continuous function* f *on* $[0, c_r/r]$ *one has*

$$\lim_{n\to\infty} \frac{1}{rn} \sum_{k=1}^{rn} f\left(\frac{x_{k,rn}}{rn}\right) = \int_0^{c_r} f(t/r) u_r(t)\, dt, \qquad c_r = \frac{(r+1)^{r+1}}{r^r},$$

where the density u_r *on* $[0, c_r]$ *is given by*

$$u_r(\hat{x}) = \frac{1}{r\pi} \frac{(\sin r\varphi)^{r+1}}{\big(\sin(r+1)\varphi\big)^r}, \qquad 0 < \varphi < \frac{\pi}{r+1}, \tag{1.5}$$

where

$$\hat{x} = \frac{\big(\sin(r+1)\varphi\big)^{r+1}}{\sin\varphi\big(\sin r\varphi\big)^r}, \qquad 0 < \varphi < \frac{\pi}{r+1}.$$

The densities u_r for $1 \le r \le 5$ are plotted in Figure 2.



Figure 2: The asymptotic zero densities u_r for multiple Laguerre polynomials of the first kind: $r = 1$ (solid), $r = 2$ (dash), $r = 3$ (dash dot), $r = 4$ (long dash), and $r = 5$ (dots).

The density of the scaled zeros $\{\frac{x_{k,rn}}{rn}, 1 \le k \le rn\}$ is therefore given by $ru_r(rx)$ for $0 < x < c_r/r$. Note that the densities u_r behave as

$$u_r(\hat{x}) = \mathcal{O}(\varphi) = \mathcal{O}\big((c_r - \hat{x})^{1/2}\big), \qquad \hat{x} \to c_r,$$

and

$$u_r(\hat{x}) = \mathcal{O}\left(\left(\frac{\pi}{r+1} - \varphi\right)^{-r}\right) = \mathcal{O}\left(\hat{x}^{-\frac{r}{r+1}}\right), \qquad \hat{x} \to 0.$$

Hence the densities u_r tend to zero as a square root near the endpoint c_r and have the same singularity near 0 as in the Jacobi-Piñeiro case. For $r = 1$ the density is the

Marchenko-Pastur density [12]

$$u_1(\hat{x}) = \frac{1}{2\pi}\sqrt{\frac{4-\hat{x}}{\hat{x}}}, \qquad 0 < \hat{x} < 4, \tag{1.6}$$

which is also the known asymptotic distribution of the (scaled) zeros of Laguerre polynomials (see, e.g., [6]). For $r = 2$ we have $u_2(\hat{x}) = \frac{8}{27} g(\frac{8\hat{x}}{27})$, where

$$g(y) = \frac{3\sqrt{3}}{16\pi} \frac{(1+3\sqrt{1-y})(1-\sqrt{1-y})^{1/3} - (1-3\sqrt{1-y})(1+\sqrt{1-y})^{1/3}}{y^{2/3}},$$

and the asymptotic zero distribution of the zeros of multiple Laguerre polynomials for that case was already obtained in [3, Thm. 2.6]. An interesting observation is that the moments of u_r are given by

$$\int_0^{c_r} x^n u_r(x)\, dx = \frac{1}{n+1}\binom{(r+1)n}{n}, \qquad n \in \mathbb{N}.$$

The simple expressions for the moments of w_r and u_r on $[0, c_r]$ is the main reason why we prefer to express the asymptotic zero densities in terms of densities on $[0, c_r]$, rather than on $[0, 1]$ and $[0, c_r/r]$ respectively. In Section 3 we will show that these densities and the asymptotic behavior of the ratio of Jacobi-Piñeiro and multiple Laguerre polynomials of the first kind are related to the Fuss-Catalan distribution with density

$$g_r(x) = \frac{1}{\pi} \frac{\sin^2\varphi \big(\sin r\varphi\big)^{r-1}}{\big(\sin(r+1)\varphi\big)^r}, \qquad 0 < x < c_r,$$

where

$$x = \frac{\big(\sin(r+1)\varphi\big)^{r+1}}{\sin\varphi \big(\sin r\varphi\big)^r}, \qquad 0 < \varphi < \frac{\pi}{r+1},$$

for which the moments are the Fuss-Catalan numbers [7, p. 347]

$$\int_0^{c_r} x^n g_r(x)\, dx = \frac{1}{rn+1}\binom{(r+1)n}{n}, \qquad n \in \mathbb{N}.$$

2 Ratio asymptotics for Jacobi-Piñeiro polynomials

The nearest neighbor recurrence relations are

$$xP_{\vec{n}}(x) = P_{\vec{n}+\vec{e}_k}(x) + b_{\vec{n},k}P_{\vec{n}}(x) + \sum_{j=1}^r a_{\vec{n},j}P_{\vec{n}-\vec{e}_j}(x), \qquad 1 \le k \le r,$$

where the recurrence coefficients are given by

$$\begin{aligned} a_{\vec{n},j} = {} & \frac{n_j(n_j+\alpha_j)(|\vec{n}|+\beta)}{(|\vec{n}|+n_j+\alpha_j+\beta+1)(|\vec{n}|+n_j+\alpha_j+\beta)(|\vec{n}|+n_j+\alpha_j+\beta-1)} \\ & \times \prod_{i=1}^r \frac{|\vec{n}|+\alpha_i+\beta}{|\vec{n}|+n_i+\alpha_i+\beta} \prod_{i=1, i\neq j}^r \frac{n_j+\alpha_j-\alpha_i}{n_j-n_i+\alpha_j-\alpha_i}, \qquad 1 \le j \le r, \end{aligned} \tag{2.1}$$

and

$$b_{\vec{n},k} = (|\vec{n}|+\beta+1)\frac{\prod_{j=1}^{r}(|\vec{n}|+\beta+\alpha_j+1)}{(|\vec{n}|+n_k+\beta+\alpha_k+2)\prod_{j\neq k}(|\vec{n}|+n_j+\beta+\alpha_j+1)} - (|\vec{n}|+\beta)\frac{\prod_{j=1}^{r}(|\vec{n}|+\beta+\alpha_j)}{\prod_{j=1}^{r}(|\vec{n}|+n_j+\beta+\alpha_j)}, \qquad 1 \le k \le r. \tag{2.2}$$

(see, e.g., [20]).

If we take the multi-index $\vec{n} = (\lfloor q_1 n\rfloor, \ldots, \lfloor q_r n\rfloor)$, where $q_j > 0$, $\sum_{j=1}^{r} q_j = 1$, and $\lfloor\cdot\rfloor$ is the floor function (i.e., $\lfloor a\rfloor = k$ whenever $k \le a < k+1$), then the asymptotic behavior of the recurrence coefficients is

$$\lim_{n\to\infty} a_{\vec{n},j} = \frac{q_j^{r+1}}{(1+q_j)^3}\prod_{k=1}^{r}\frac{1}{1+q_k}\prod_{i\neq j}\frac{1}{q_j-q_i} =: a_j, \qquad 1 \le j \le r, \tag{2.3}$$

and with a bit of elementary calculus

$$\lim_{n\to\infty} b_{\vec{n},j} = \prod_{k=1}^{r}\frac{1}{1+q_k}\left(r+1-\sum_{k=1}^{r}\frac{1}{1+q_k}-\frac{1}{1+q_j}\right) =: b_j, \qquad 1 \le j \le r. \tag{2.4}$$

In order to have finite values of a_j, we assume for the moment that $q_i \neq q_j$ whenever $i \neq j$, but later on we will take the limit $q_j \to 1/r$ for every j. This passage to the limit is allowed since the asymptotic distribution of the zeros is continuous in the parameters $(q_1, \ldots, q_r)$, which can be shown as in [4, Thm. 2]. We will use the notation

$$p(\vec{q}) = \prod_{k=1}^{r}\frac{1}{1+q_k}, \quad s = r+1-\sum_{k=1}^{r}\frac{1}{1+q_k},$$

so that

$$a_j = p(\vec{q})\frac{q_j^{r+1}}{(1+q_j)^3}\prod_{i\neq j}\frac{1}{q_j-q_i}, \quad b_j = p(\vec{q})\left(s-\frac{1}{1+q_j}\right). \tag{2.5}$$

According to [21, Thm. 1.1], the ratio asymptotics for the Jacobi-Piñeiro polynomials with multi-index $\vec{n} = (\lfloor q_1 n\rfloor, \ldots, \lfloor q_r n\rfloor)$ will then be given by

$$\lim_{n\to\infty}\frac{P_{\vec{n}+\vec{e}_k}(x)}{P_{\vec{n}}(x)} = z(x) - b_k, \qquad 1 \le k \le r, \tag{2.6}$$

uniformly on compact subsets of $\mathbb{C}\setminus[0,1]$, where z is the solution of the algebraic equation

$$(z-x)B_r(z) + A_{r-1}(z) = 0 \tag{2.7}$$

for which $z(x) - x \to 0$ when $x \to \infty$. In [21] the convergence was given uniformly on compact subsets of $\mathbb{C}\setminus\mathbb{R}$, but since all the zeros of Jacobi-Piñeiro polynomials are in $[0,1]$, the Stieltjes-Vitali theorem can be used to extend the uniform convergence to compact subsets of $\mathbb{C}\setminus[0,1]$. Here $B_r(z) = \prod_{j=1}^{r}(z-b_j)$ and A_{r-1} is the polynomial of degree $r-1$ for which

$$\frac{A_{r-1}(z)}{B_r(z)} = \sum_{j=1}^{r}\frac{a_j}{z-b_j}.$$

The latter means that the residue of A_{r-1}/B_r at b_j is given by a_j:

$$a_j = \frac{A_{r-1}(b_j)}{B_r'(b_j)} = \frac{A_{r-1}(b_j)}{\prod_{i\neq j}(b_j - b_i)}. \tag{2.8}$$

Observe that

$$\prod_{i\neq j}(b_j - b_i) = \frac{p(\vec{q})^r}{(1+q_j)^{r-2}} \prod_{i\neq j}(q_j - q_i),$$

so that the condition on the residues (2.8) becomes

$$A_{r-1}(b_j) = \left(\frac{p(\vec{q})q_j}{1+q_j}\right)^{r+1}, \qquad 1 \leq j \leq r. \tag{2.9}$$

This is a Lagrange interpolation problem. If we use (2.5) to write q_j in terms of b_j, then

$$q_j = \frac{1}{s - b_j/p(\vec{q})} - 1,$$

so that

$$\frac{p(\vec{q})q_j}{1+q_j} = p(\vec{q})(1-s) + b_j, \qquad 1 \leq j \leq r.$$

The interpolation problem (2.9) then becomes

$$A_{r-1}(b_j) = \big(b_j + p(\vec{q})(1-s)\big)^{r+1}, \qquad 1 \leq j \leq r,$$

hence $A_{r-1}(z)$ is a polynomial of degree $r-1$ interpolating the polynomial $\big(z + p(\vec{q})(1-s)\big)^{r+1}$ at the points b_j $(1 \leq j \leq r)$. If we take the limit where $q_j \to 1/r$ for every j, then

$$p(\vec{q}) \to \left(\frac{r}{r+1}\right)^r =: p, \quad s \to \frac{2r+1}{r+1}, \quad b_j \to p\left(s - \frac{r}{r+1}\right) = p,$$

hence all the interpolation points coincide. It is well known that the Lagrange interpolating polynomial for which all the interpolation points coincide corresponds to the Taylor polynomial of degree $r-1$ of the function $f(z) = \big(z + p(1-s)\big)^{r+1}$ around the common interpolation point p. This Taylor polynomial of degree $r-1$ is the polynomial $\big(z+p(1-s)\big)^{r+1}$ of degree $r+1$ from which we subtract the last two terms of the Taylor expansion around p:

$$\begin{aligned} A_{r-1}(z) &= \big(z+p(1-s)\big)^{r+1} - (z-p)^{r+1}\frac{f^{(r+1)}(p)}{(r+1)!} - (z-p)^r\frac{f^{(r)}(p)}{r!} \\ &= \big(z+p(1-s)\big)^{r+1} - (z-p)^{r+1} - (r+1)p(2-s)(z-p)^r. \end{aligned} \tag{2.10}$$

The algebraic equation (2.7) for multi-indices on the diagonal then becomes

$$(z-x)(z-p)^r + \big(z+p(1-s)\big)^{r+1} - (z-p)^{r+1} - (r+1)p(2-s)(z-p)^r = 0,$$

which simplifies to

$$x(z-p)^r = \left(z - \frac{pr}{r+1}\right)^{r+1}. \tag{2.11}$$

3 Relation with the Fuss-Catalan numbers

Recently the Fuss-Catalan distribution and other related distributions (Raney distributions) appeared as limiting distributions of eigenvalues and singular values of certain random matrices [5], [15], [16]. In this section we will show how the ratio asymptotics in (2.6) is related to the Stieltjes transform of the Fuss-Catalan distribution. The weights w_r and u_r in Theorem 1.1 and 1.2 cannot be identified with the Fuss-Catalan distribution or any of the Raney distributions (except u_1, which is the Catalan distribution) because their behavior near the endpoints of the interval differs from the behavior of the Raney distributions given in [13].

If we scale the variables $\hat{x} = c_r x$ and $\hat{z} = c_r z$, where

$$c_r = \frac{(r+1)^{r+1}}{r^r} = \frac{r+1}{p},$$

then the algebraic equation (2.11) becomes

$$\hat{x}(\hat{z} - r - 1)^r = (\hat{z} - r)^{r+1}. \tag{3.1}$$

If we define

$$\omega = \frac{\hat{z} - r}{\hat{z} - r - 1}, \quad \hat{z} = \frac{(r+1)\omega - r}{\omega - 1}, \tag{3.2}$$

then the algebraic equation becomes

$$\omega^{r+1} + \hat{x} - \hat{x}\omega = 0. \tag{3.3}$$

This is the algebraic equation for the generating function $G(1/\hat{x})$ of the Fuss-Catalan numbers [7, p. 347] [16, Eq. (3.12)]. As in [16, §3], we assume that a solution exists of the form $\omega = \rho e^{i\varphi}$, where $\rho > 0$ and φ is real. Then inserting this in (3.3) gives

$$\rho^{r+1} e^{i(r+1)\varphi} + \hat{x} - \hat{x}\rho e^{i\varphi} = 0.$$

This gives for the real and the imaginary part

$$\begin{aligned} \rho^{r+1}\cos(r+1)\varphi + \hat{x} - \hat{x}\rho\cos\varphi &= 0, & (3.4) \\ \rho^{r+1}\sin(r+1)\varphi - \hat{x}\rho\sin\varphi &= 0. & (3.5) \end{aligned}$$

From (3.5) we find

$$\hat{x} = \rho^r \frac{\sin(r+1)\varphi}{\sin\varphi}, \tag{3.6}$$

and inserting this in (3.4) gives

$$\rho(\hat{x}) = \frac{\sin(r+1)\varphi}{\sin r\varphi}, \tag{3.7}$$

from which

$$\hat{x} = \frac{\big(\sin(r+1)\varphi\big)^{r+1}}{\sin\varphi\big(\sin r\varphi\big)^r}. \tag{3.8}$$

Observe that $\rho(x) > 0$ for $0 < \varphi < \frac{\pi}{r+1}$, and $\hat{x}$ is a monotonically decreasing function mapping $[0, \frac{\pi}{r+1}]$ into $[0, c_r]$. So for $\hat{x} \in [0, c_r]$ there is a solution $\rho e^{i\varphi}$ of the algebraic equation (3.3). The conjugate function $\rho e^{-i\varphi}$ is also a solution for $\hat{x} \in [0, c_r]$. In fact both solutions are the boundary value of a function ω which is analytic on $\mathbb{C} \setminus [0, c_r]$ and

$$\omega_+ = \lim_{\epsilon \to 0+} \omega(\hat{x} + i\epsilon) = \rho e^{-i\phi}, \quad \omega_- = \lim_{\epsilon \to 0+} \omega(\hat{x} - i\epsilon) = \rho e^{i\phi}, \tag{3.9}$$

because this ω is $G(1/\hat{x}) = \hat{x}F(\hat{x})$, where F is the Stieltjes transform of the Fuss-Catalan distribution

$$F(z) = \int_0^{c_r} \frac{g_r(y)}{z - y}\, dy, \qquad z \in \mathbb{C} \setminus [0, c_r],$$

with g_r the Fuss-Catalan density, and a Stieltjes transform has the property that

$$\operatorname{Im} F(z) \begin{cases} < 0, & \operatorname{Im} z > 0, \\ > 0, & \operatorname{Im} z < 0. \end{cases}.$$

Observe that

$$\frac{1}{z - p} = \lim_{n \to \infty} \frac{P_{\vec{n}}(x)}{P_{\vec{n}+\vec{e}_k}(x)}$$

is the Stieltjes transform of a probability measure on $[0, 1]$, since we have

$$\frac{P_{\vec{n}}(x)}{P_{\vec{n}+\vec{e}_k}(x)} = \sum_{j=1}^{|\vec{n}|+1} \frac{c_{j,\vec{n}}}{x - x_{j,\vec{n}+\vec{e}_k}},$$

and $c_{j,\vec{n}} > 0$ because the zeros of $P_{\vec{n}}$ and $P_{\vec{n}+\vec{e}_k}$ interlace [8, Thm.2.1], and $\sum c_{j,\vec{n}} = 1$ since we are dealing with monic polynomials. With the change of variables $\hat{x} = c_r x$ and $\hat{z} = c_r z$ it follows that

$$1/(\hat{z} - r - 1) = \int_0^{c_r} \frac{d\mu(y)}{\hat{x} - y}$$

is the Stieltjes transform of a probability distribution μ on $[0, c_r]$. Note that (3.2) implies

$$\frac{1}{\hat{z} - r - 1} = \omega - 1 = \hat{x}F(\hat{x}) - 1,$$

where F is the Stieltjes transform of the Fuss-Catalan distribution,

$$F(\hat{x}) = \frac{1}{\hat{x}} \sum_{n=0}^{\infty} \frac{1}{rn + 1} \binom{(r+1)n}{n} \frac{1}{\hat{x}^n},$$

so that $1/(\hat{z} - r - 1)$ is the Stieltjes transform of the probability measure for which the moments are the Fuss-Catalan numbers shifted by one

$$\int_0^{c_r} y^n\, d\mu(y) = \frac{1}{r(n+1) + 1} \binom{(r+1)(n+1)}{n+1},$$

and hence this probability distribution has a density $\hat{x}g_r(\hat{x})$, where g_r is the Fuss-Catalan density

$$g_r(\hat{x}) = \frac{1}{\pi} \frac{\sin^2 \varphi (\sin r\varphi)^{r-1}}{\big(\sin(r+1)\varphi\big)^r}, \qquad 0 < \varphi < \frac{\pi}{r+1},$$

where $\hat{x}$ is given in (3.8). In particular this gives

$$\frac{1}{\hat{z}-r-1} = \int_0^{c_r} \frac{y g_r(y)}{\hat{x}-y}\, dy.$$

The weight is explicitly given by

$$\hat{x} g_r(\hat{x}) = \frac{1}{\pi} \frac{\sin\varphi \sin(r+1)\varphi}{\sin r\varphi}, \qquad 0 \leq \varphi < \frac{\pi}{r+1},$$

with $\hat{x}$ as in (3.8).

4 Proof of Theorem 1.1

So far we found that for $\vec{n}$ near the diagonal (i.e., $n_j/n \to 1/r$ for every j) one has

$$\lim_{n\to\infty} \frac{P_{\vec{n}+\vec{e}_k}(x)}{P_{\vec{n}}(x)} = z(x) - p = \frac{1}{c_r}(\hat{z}-r-1), \tag{4.1}$$

uniformly for x on compact subsets of $\mathbb{C} \setminus [0,1]$, or $\hat{x}$ on compact subsets of $\mathbb{C} \setminus [0,c_r]$. However we are interested in the asymptotic behavior of

$$\frac{1}{|\vec{n}|} \frac{P'_{\vec{n}}(x)}{P_{\vec{n}}(x)},$$

where the prime $'$ denotes the derivative with respect to x, because the limit will give the Stieltjes transform of the asymptotic distribution of the zeros of $P_{\vec{n}}$. By taking derivatives with respect to x in (4.1) we find

$$\lim_{n\to\infty} \frac{P_{\vec{n}+\vec{e}_k}(x)}{P_{\vec{n}}(x)} \left(\frac{P'_{\vec{n}+\vec{e}_k}(x)}{P_{\vec{n}+\vec{e}_k}(x)} - \frac{P'_{\vec{n}}(x)}{P_{\vec{n}}(x)} \right) = z' = \frac{\hat{z}'}{c_r},$$

uniformly for x on compact subsets of $\mathbb{C} \setminus [0,1]$. Together with (4.1), this gives

$$\lim_{n\to\infty} \left(\frac{P'_{\vec{n}+\vec{e}_k}(x)}{P_{\vec{n}+\vec{e}_k}(x)} - \frac{P'_{\vec{n}}(x)}{P_{\vec{n}}(x)} \right) = \frac{\hat{z}'}{\hat{z}-r-1}.$$

If we use this result successively for each k, $1 \leq k \leq r$, then we find for multi-indices on the diagonal $n\vec{1} = (n,n,\ldots,n)$ and $(n+1)\vec{1} = (n+1,n+1,\ldots,n+1)$

$$\lim_{n\to\infty} \left(\frac{P'_{(n+1)\vec{1}}(x)}{P_{(n+1)\vec{1}}(x)} - \frac{P'_{n\vec{1}}(x)}{P_{n\vec{1}}(x)} \right) = \frac{r\hat{z}'}{\hat{z}-r-1}.$$

Then by taking averages (Cesàro's lemma) we get

$$\lim_{n\to\infty} \frac{1}{n} \sum_{k=0}^{n-1} \left(\frac{P'_{(k+1)\vec{1}}(x)}{P_{(k+1)\vec{1}}(x)} - \frac{P'_{k\vec{1}}(x)}{P_{k\vec{1}}(x)} \right) = \frac{r\hat{z}'}{\hat{z}-r-1},$$

and since this contains a telescoping sum, this becomes

$$\lim_{n\to\infty} \frac{1}{rn} \frac{P'_{n\vec{1}}(x)}{P_{n\vec{1}}(x)} = \frac{\hat{z}'}{\hat{z}-r-1},$$

uniformly for x on compact subsets of $\mathbb{C} \setminus [0,1]$, so that the right hand side is the Stieltjes transform of the asymptotic zero distribution of the zeros of $P_{n\vec{1}}$ for $n\vec{1} = (n, n, \dots, n)$. From the Stieltjes transform we can find the density by using Stieltjes' inversion formula

$$2\pi i v_r(x) = \left(\frac{\hat{z}'}{\hat{z} - r - 1} \right)_- - \left(\frac{\hat{z}'}{\hat{z} - r - 1} \right)_+ .$$

Taking derivatives in (3.1) (and recalling that $\hat{x} = c_r x$) gives

$$c_r (\hat{z} - r - 1)^r + \hat{x} r (\hat{z} - r - 1)^{r-1} \hat{z}' = (r+1)\,(\hat{z} - r)^r\, \hat{z}',$$

so that

$$\frac{\hat{z}'}{\hat{z} - r - 1} = \frac{c_r}{\hat{x}} \frac{\hat{z} - r}{\hat{z} - 2r - 1}.$$

Writing this in terms of ω using (3.2) gives

$$\frac{\hat{z}'}{\hat{z} - r - 1} = \frac{c_r}{\hat{x}} \frac{\omega}{-r\omega + r + 1}.$$

Now use $\omega_+ = \rho e^{-i\varphi}$ and $\omega_- = \rho e^{i\varphi}$ to find the density

$$v_r(x) = \frac{r+1}{\pi x} \frac{\rho \sin \varphi}{|r\rho e^{i\varphi} - r - 1|^2},$$

and clearly $v_r(x) = v_r(\hat{x}/c_r) = c_r w_r(\hat{x})$, with the weight in (1.4). Observe that $\hat{x} : [0, \frac{\pi}{r+1}] \to [0, c_r]$ is a monotonically decreasing function with

$$\hat{x}'(\varphi) = \frac{-\hat{x}}{\sin \varphi \sin r\varphi \sin(r+1)\varphi} |(r+1) \sin r\varphi - e^{i\varphi} r \sin(r+1)\varphi|^2$$

so that

$$w_r(\hat{x}) = \frac{r+1}{\pi} \frac{1}{|\hat{x}'(\varphi)|}, \qquad 0 < \varphi < \frac{\pi}{r+1}.$$

5 Ratio asymptotics for multiple Laguerre polynomials of the first kind

The nearest neighbor recurrence relations for multiple orthogonal polynomials of the first kind are given by

$$x L_{\vec{n}}(x) = L_{\vec{n}+\vec{e}_k}(x) + b_{\vec{n},k} L_{\vec{n}}(x) + \sum_{j=1}^{r} a_{\vec{n},j} L_{\vec{n}-\vec{e}_j}(x), \qquad 1 \le k \le r,$$

where the recurrence coefficients are given by

$$a_{\vec{n},j} = n_j (n_j + \alpha_j) \prod_{i=1, i \neq j}^{r} \frac{n_j + \alpha_j - \alpha_i}{n_j - n_i + \alpha_j - \alpha_i}, \qquad 1 \le j \le r, \tag{5.1}$$

and

$$b_{\vec{n},k} = |\vec{n}| + n_k + \alpha_k + 1, \qquad 1 \le k \le r. \tag{5.2}$$

(see, e.g., [20]). We can now proceed as in the case of Jacobi-Piñeiro polynomials. The recurrence coefficients are somewhat easier but they are unbounded so that we need to use a scaling. Suppose again that $\vec{n} = (\lfloor q_1 n \rfloor, \ldots, \lfloor q_r n \rfloor)$, where $q_i \neq q_j$ whenever $i \neq j$. It then follows that

$$\lim_{n\to\infty} \frac{a_{\vec{n},j}}{n^2} = q_j^{r+1} \prod_{i\neq j} \frac{1}{q_j - q_i} =: a_j, \qquad 1 \leq j \leq r,$$

and

$$\lim_{n\to\infty} \frac{b_{\vec{n},j}}{n} = 1 + q_j =: b_j, \qquad 1 \leq j \leq r.$$

According to [21, Thm. 1.2] we then have

$$\lim_{n\to\infty} \frac{L_{\vec{n}+\vec{e}_k}(nx)}{nL_{\vec{n}}(nx)} = z(x) - b_k, \qquad 1 \leq k \leq r, \tag{5.3}$$

uniformly on compact subsets of $\mathbb{C}\backslash[0,\infty)$, where z is the solution of the algebraic equation

$$(z-x)B_r(z) + A_{r-1}(z) = 0,$$

where $B_r(z) = \prod_{j=1}^r (z - b_j)$ and A_{r-1} is obtained from

$$\frac{A_{r-1}(z)}{B_r(z)} = \sum_{j=1}^r \frac{a_j}{z - b_j}.$$

The uniform convergence on compact subsets of $\mathbb{C}\setminus\mathbb{R}$ in [21] can be extended to $\mathbb{C}\setminus[0,\infty)$ because the zeros of multiple Laguerre polynomials of the first kind are on $[0,\infty)$. One can even extend this further to $\mathbb{C}\setminus[0, c_r/r]$ since all the scaled zeros are dense on $[0, c_r/r]$, but we will not need this here. Observe that

$$\prod_{i\neq j}(q_j - q_i) = \prod_{i\neq j}(b_j - b_i),$$

so that we get the interpolation condition

$$A_{r-1}(b_j) = q_j^{r+1} = (b_j - 1)^{r+1}, \qquad 1 \leq j \leq r.$$

Hence $A_{r-1}(z)$ is the Lagrange interpolating polynomial of degree $r-1$ for the function $f(z) = (z-1)^{r+1}$ for the interpolation points $b_1, \ldots, b_r$. Now let $q_j \to 1/r$ for every j, then

$$b_j \to \frac{r+1}{r},$$

and $A_{r-1}(z)$ will be the Taylor polynomial of degree $r-1$ around $\frac{r+1}{r}$ for the function $f(z) = (z-1)^{r+1}$. This gives

$$A_{r-1}(z) = (z-1)^{r+1} - \left(z - \frac{r+1}{r}\right)^{r+1} - \frac{r+1}{r}\left(z - \frac{r+1}{r}\right)^r.$$

The algebraic equation for multi-indices near the diagonal then becomes

$$x\left(z - \frac{r+1}{r}\right)^r = (z-1)^{r+1}. \tag{5.4}$$

The change of variables $rz = \hat{z}$ and $rx = \hat{x}$ gives the same algebraic equation as in (3.1).

6 Proof of Theorem 1.2

As in Section 4 we use

$$\frac{L'_{n\vec{1}}(x)}{L_{n\vec{1}}(x)} = \sum_{k=0}^{n-1} \left(\frac{L'_{(k+1)\vec{1}}(x)}{L_{(k+1)\vec{1}}(x)} - \frac{L'_{k\vec{1}}(x)}{L_{k\vec{1}}(x)} \right),$$

where $k\vec{1} = (k,k,\ldots,k)$ and $(k+1)\vec{1} = (k+1,k+1,\ldots,k+1)$. However, because of the scaling, we need to consider (observe that $|n\vec{1}| = rn$)

$$\frac{L'_{n\vec{1}}(rnx)}{rnL_{n\vec{1}}(rnx)} = \sum_{k=0}^{n-1} \left(\frac{L'_{(k+1)\vec{1}}(rnx)}{rnL_{(k+1)\vec{1}}(rnx)} - \frac{L'_{k\vec{1}}(rnx)}{rnL_{k\vec{1}}(rnx)} \right),$$

so that we can not use Cesàro's lemma to get the asymptotic behavior. We modify the proof as follows. For $\frac{k}{n} \leq t < \frac{k+1}{n}$ one has $\lfloor nt \rfloor = k$, hence the sum can be written as an integral

$$\frac{L'_{n\vec{1}}(rnx)}{rnL_{n\vec{1}}(rnx)} = n \int_0^1 \left(\frac{L'_{(\lfloor nt \rfloor+1)\vec{1}}(rnx)}{rnL_{(\lfloor nt \rfloor+1)\vec{1}}(rnx)} - \frac{L'_{\lfloor nt \rfloor\vec{1}}(rnx)}{rnL_{\lfloor nt \rfloor\vec{1}}(rnx)} \right) dt,$$

and the integrand can be written as

$$\left(\frac{L_{(\lfloor nt \rfloor+1)\vec{1}}(rnx)}{rnL_{\lfloor nt \rfloor\vec{1}}(rnx)} \right)' \Big/ \left(\frac{L_{(\lfloor nt \rfloor+1)\vec{1}}(rnx)}{rnL_{\lfloor nt \rfloor\vec{1}}(rnx)} \right).$$

So we need to know the asymptotic behavior of the ratio

$$\lim_{n\to\infty} \frac{L_{(\lfloor nt \rfloor+1)\vec{1}}(rnx)}{rnL_{\lfloor nt \rfloor\vec{1}}(rnx)}.$$

If we change n to rn in Section 5 then for $q_j \to \frac{1}{r}$ $(1 \leq j \leq r)$ we get the multi-index $n\vec{1} = (n,n,\ldots,n)$ and (5.3) becomes

$$\lim_{n\to\infty} \frac{L_{n\vec{1}+\vec{e}_k}(rnx)}{rnL_{n\vec{1}}(rnx)} = z(x) - \frac{r+1}{r},$$

but we need to extend this for multi-indices containing the parameter $0 < t \leq 1$. For this we need to use the following asymptotic behavior of the recurrence coefficients: if $\vec{n} = (\lfloor nq_1 \rfloor, \lfloor nq_2 \rfloor, \ldots, \lfloor nq_r \rfloor)$ and $\vec{m} = (\lfloor ntq_1 \rfloor, \lfloor ntq_2 \rfloor, \ldots, \lfloor ntq_r \rfloor)$, then

$$\lim_{n\to\infty} \frac{a_{\vec{m},j}}{n^2} = t^2 q_j^{r+1} \prod_{i\neq j} \frac{1}{q_j - q_i} = t^2 a_j, \qquad 1 \leq j \leq r,$$

and

$$\lim_{n\to\infty} \frac{b_{\vec{m},j}}{n} = t(1+q_j) = tb_j, \qquad 1 \leq j \leq r.$$

The required asymptotic behavior is then for $0 < t \leq 1$

$$\lim_{n\to\infty} \frac{L_{\vec{m}+\vec{e}_k}(nx)}{nL_{\vec{m}}(nx)} = z(x,t) - tb_k, \tag{6.1}$$

uniformly for x on compact subsets of $\mathbb{C} \setminus [0,\infty)$, where $z(x,t)$ satisfies the algebraic equation

$$\big(z(x,t) - x\big) B_r(z,t) + A_{r-1}(z,t) = 0,$$

with $B_r(z,t) = \prod_{j=1}^r (z - tb_j) = t^r B_r(z/t)$ and

$$\frac{A_{r-1}(z,t)}{B_r(z,t)} = \sum_{j=1}^r \frac{t^2 a_j}{z - tb_j},$$

so that $A_{r-1}(z,t) = t^{r+1} A_{r-1}(z/t)$. Here we used $A_{r-1}(z) = A_{r-1}(z,1)$ and $B_r(z) = B_r(z,1)$, which are the polynomials in Section 5. If $q_j \to \frac{1}{r}$ $(1 \leq j \leq r)$ then $b_j \to \frac{r+1}{r}$ $(1 \leq j \leq r)$ and the algebraic equation for $z(x,t)$ becomes

$$x \left(z(x,t) - t\frac{r+1}{r} \right)^r = \big(z(x,t) - t\big)^{r+1}. \tag{6.2}$$

Now change n to rn so that we can deal with the multi-index $n\vec{1} = (n, n, \ldots, n)$. By going from the multi-index $n\vec{1} = (n, n, \ldots, n)$ to $(n+1)\vec{1} = (n+1, n+1, \ldots, n+1)$ in r steps (each time increasing one coefficient) we then get

$$\lim_{n\to\infty} \frac{L_{(\lfloor nt \rfloor + 1)\vec{1}}(rnx)}{(rn)^r L_{\lfloor nt \rfloor \vec{1}}(rnx)} = \left(z(x,t) - t\frac{r+1}{r} \right)^r,$$

so that

$$\lim_{n\to\infty} \frac{1}{rn} \frac{L'_{n\vec{1}}(rnx)}{rnL_{n\vec{1}}(rnx)} = \frac{1}{r} \int_0^1 \frac{\frac{d}{dx}\left(z(x,t) - t\frac{r+1}{r}\right)^r}{\left(z(x,t) - t\frac{r+1}{r}\right)^r}\, dt = \int_0^1 \frac{z'(x,t)}{z(x,t) - t\frac{r+1}{r}}\, dt,$$

uniformly on compact subsets of $\mathbb{C} \setminus [0,\infty)$, where the prime $'$ means the derivative with respect to x. This limit is the Stieltjes transform of the asymptotic zero distribution

$$\int_0^{c_r/r} r\frac{u_r(rs)}{x - s}\, ds = \int_0^{c_r} \frac{u_r(y)}{x - y/r}\, dy,$$

and hence we have

$$\int_0^1 \frac{z'(x,t)}{z(x,t) - t\frac{r+1}{r}}\, dt = \int_0^{c_r} \frac{u_r(y)}{x - y/r}\, dy.$$

Observe that the change of variables $rz = t\hat{z}$ and $rx = t\hat{x}$ transforms the algebraic equation (6.2) to (3.1), so that $z(t\hat{x}/r, t) = t\hat{z}(\hat{x})/r$. From our analysis in Sections 2–4 we found that

$$\frac{\hat{z}'}{\hat{z} - r - 1} = \int_0^{c_r} \frac{w_r(s)}{\hat{x} - s}\, ds,$$

hence

$$\frac{z'(x,t)}{z(x,t) - t\frac{r+1}{r}} = \frac{r}{t} \int_0^{c_r} \frac{w_r(s)}{\frac{rx}{t} - s}\, ds = \int_0^{c_r} \frac{w_r(s)}{x - \frac{ts}{r}}\, ds.$$

Therefore

$$\begin{aligned}
\int_0^1 \frac{z'(x,t)}{z(x,t) - t\frac{r+1}{r}}\, dt &= \int_0^1 \int_0^{c_r} \frac{w_r(s)}{x - \frac{ts}{r}}\, ds\, dt \\
&= \int_0^1 \int_0^{tc_r} \frac{w_r(y/t)}{x - \frac{y}{r}} \frac{dy}{t}\, dt \\
&= \int_0^{c_r} \frac{1}{x - \frac{y}{r}} \int_{y/c_r}^1 w_r(y/t) \frac{dt}{t}\, dy,
\end{aligned}$$

where we used the change of variables $ts = y$ in the second equality and Fubini's theorem for the third equality. This means that

$$u_r(y) = \int_{y/c_r}^{1} w_r(y/t)\,\frac{dt}{t} = \int_{y}^{c_r} w_r(x)\,\frac{dx}{x}, \tag{6.3}$$

and hence the asymptotic density of the scaled zeros u_r is the Mellin convolution of the density w_r given in (1.4) and the uniform distribution on $[0,1]$. This immediately gives the moments

$$\int_0^{c_r} y^n u_r(y)\,dy = \int_0^{c_r} x^n w_r(x)\,dx \int_0^1 t^n\,dt = \frac{1}{n+1}\binom{(r+1)n}{n}.$$

We still need to show that the density u_r is given by the expression in (1.5). Observe that the derivative of (1.5) with respect to φ is

$$\frac{r+1}{\pi}\frac{\sin\varphi(\sin r\varphi)^r}{\big(\sin(r+1)\varphi\big)^{r+1}} = \frac{r+1}{\pi\hat{x}}.$$

On the other hand, taking the derivative in (6.3) with respect to φ gives

$$\frac{du_r(\hat{x})}{d\varphi} = -\frac{w_r(\hat{x})}{\hat{x}}\hat{x}' = \frac{r+1}{\pi\hat{x}},$$

where we used (1.4) for the last equality. Thus, using $u_r(c_r) = 0$, we find that

$$u_r(\hat{x}) = \frac{1}{r\pi}\frac{(\sin r\varphi)^{r+1}}{\big(\sin(r+1)\varphi\big)^r}.$$

7 Concluding remarks

There is yet another family of multiple orthogonal polynomials for which the asymptotic distribution of the zeros is of the same flavor. These are multiple orthogonal polynomials associated with Meijer G-functions, which appear in the study of products of Ginibre random matrices [11]. The polynomials on the stepline for $|\vec{n}| = n$ are given by

$$P_n(x) = (-1)^n \prod_{j=1}^{r}(n+\nu_j)!\,\sum_{k=0}^{n}\binom{n}{k}\frac{(-x)^k}{(k+\nu_1)!\cdots(k+\nu_r)!},$$

and the asymptotic distribution of the scaled zeros $\{x_{k,n}/n^r, 1 \le k \le n\}$ is given in [16, Thm. 3.2]. The density is

$$g_r(x) = \frac{1}{\pi}\frac{\sin^2\varphi(\sin r\varphi)^{r-1}}{\big(\sin(r+1)\varphi\big)^r},$$

where again

$$x = \frac{\big(\sin(r+1)\varphi\big)^{r+1}}{\sin\varphi(\sin r\varphi)^r}, \qquad 0 < \varphi < \frac{\pi}{r+1}. \tag{7.1}$$

This is the density of the Fuss-Catalan distribution, for which the moments are the Fuss-Catalan numbers

$$\int_0^{c_r} x^n g_r(x)\,dx = \frac{1}{rn+1}\binom{(r+1)n}{n}, \qquad n \in \mathbb{N}.$$

Observe that the density g_r is a Mellin convolution of the density w_r in (1.4) and the beta($\frac{1}{r}$, 1) density:

$$\begin{aligned} g_r(y) &= \frac{1}{r}\int_{y/c_r}^{1} w_r(y/t)t^{1/r-1}\,\frac{dt}{t} \\ &= \frac{1}{r}\int_{y}^{c_r} w_r(x)\left(\frac{y}{x}\right)^{1/r-1}\frac{dx}{x} \\ &= y^{1/r-1}\frac{r+1}{r\pi}\int_0^\theta \frac{d\varphi}{x^{1/r}}, \end{aligned} \tag{7.2}$$

where

$$y(\theta) = \frac{\big(\sin(r+1)\theta\big)^{r+1}}{\sin\theta(\sin r\theta)^r}.$$

This can most easily be seen from

$$\frac{d}{d\varphi}\left(\frac{(\sin\varphi)^{1/r+1}}{\big(\sin(r+1)\varphi\big)^{1/r}}\right) = \frac{r+1}{r}\frac{1}{x^{1/r}},$$

with x given in (7.1), which enables a straightforward computation of the last integral in (7.2). The case $r = 1$ corresponds to the asymptotic zero distribution of Laguerre polynomials (the Marchenko-Pastur distribution (1.6)). The case $r = 2$ was obtained earlier in [3] and corresponds to multiple orthogonal polynomials for modified Bessel functions K_ν and $K_{\nu+1}$. The weight is then explicitly given by $g_2(x) = \frac{4}{27}h(\frac{4x}{27})$, where

$$h(y) = \frac{3\sqrt{3}}{4\pi}\frac{(1+\sqrt{1-y})^{1/3}-(1-\sqrt{1-y})^{1/3}}{y^{2/3}}, \qquad 0 < y < 1.$$

Acknowledgements

This research was supported by KU Leuven research grant OT/12/073, FWO research grant G.0934.13 and the Belgian Interuniversity Attraction Poles Programme P7/18. Thorsten Neuschel is a Research Associate (chargé de recherches) of FRS-FNRS (Belgian Fund for Scientific Research).

References

[1] M. Adler, P. van Moerbeke, D. Wang, *Random matrix minor processes related to percolation theory*, Random Matrices Theory Appl. **2** (2013), 1350008 (72 pages).

[2] A.I. Aptekarev, A. Branquinho, W. Van Assche, *Multiple orthogonal polynomials for classical weights* Trans. Amer. Math. Soc. **355** (2003), 3887–3914.

[3] E. Coussement, J. Coussement, W. Van Assche, *Asymptotic zero distribution for a class of multiple orthogonal polynomials*, Trans. Amer. Math. Soc. **360** (2008), no. 10, 5571–5588.

[4] D. Dominici, W. Van Assche, *Zero distribution of polynomials satisfying a differential-difference equation*, Anal. Appl. (Singap.) **12** (2014), no. 6, 635–666.

[5] P.J. Forrester, D.-Z. Liu, *Raney distributions and random matrix theory*, J. Stat. Phys. **158** (2015), 1051–1082.

[6] W. Gawronski, *On the asymptotic distribution of the zeros of Hermite, Leguerre, and Joncquière polynomials*, J. Approx. Theory **50** (1987), 214–231.

[7] R.L. Graham, D.E. Knuth, O. Patashnik, *Concrete Mathematics*, Addison-Wesley, Reading MA, 1989.

[8] M. Haneczok, W. Van Assche, *Interlacing properties of zeros of multiple orthogonal polynomials*, J. Math. Anal. Appl. **389** (2012), 429–438.

[9] M.E.H. Ismail, *Classical and Quantum Orthogonal Polynomials in One Variable*, Encyclopedia of Mathematics and its Applications **98**, Cambridge University Press, 2005.

[10] A.B.J. Kuijlaars, *Multiple orthogonal polynomial ensembles*, in 'Recent Trends in Orthogonal Polynomials and Approximation Theory', Contemp. Math. **507**, Amer. Math. Soc., Providence, RI, 2010, pp. 155-176.

[11] A.B.J. Kuijlaars, L. Zhang, *Singular values of products of Ginibre random matrices, multiple orthogonal polynomials and hard edge scaling limits*, Comm. Math. Phys. **332** (2014), no. 2, 759–781.

[12] V.A. Marchenko, L.A. Pastur, *Distribution of eigenvalues for some sets of random matrices*, Mat. Sb. **72** (114) (1967), no. 4, 507–536 (in Russian); translated in Math. USSR Sbornik **1** (1967), no. 4, 457–483.

[13] W. Młotkowski, K.A. Penson, K. Życzkowski, *Densities of the Raney distributions*, Doc. Math. **18** (2013), 1573–1596.

[14] E. Mukhin, A. Varchenko, *Multiple orthogonal polynomials and a counterexample to the Gaudin Bethe Ansatz conjecture*, Trans. Amer. Math. Soc. 359 (2007), no. 11, 5383–5418.

[15] K.A. Penson, K. Życzkowski, *Product of Ginibre matrices: Fuss-Catalan and Raney distributions*, Phys. Rev. E **83** (2011), 061118 (9 pages).

[16] T. Neuschel, *Plancherel-Rotach formulae for average characteristic polynomials of products of Ginibre random matrices and the Fuss-Catalan distribution*, Random Matrices Theory Appl. **3** (2014), no. 1, 1450003, 18 pp.

[17] L.R. Piñeiro Díaz, *On simultaneous approximations for some collection of Markov functions*, Vestnik Moskov. Univ. Ser. I Mat. Mekh. (1987), no. 2, 67–70 (in Russian); translated in Moscow Univ. Math. Bull. **42** (2) (1987), 52–55.

[18] G. Szegő, *Orthogonal Polynomials*, Amer. Math. Soc. Colloq. Publ. **23**, Providence RI, 1939 (fourth edition 1975).

[19] W. Van Assche, E. Coussement, *Some classical multiple orthogonal polynomials*, Numerical analysis 2000, Vol. V, Quadrature and orthogonal polynomials. J. Comput. Appl. Math. **127** (2001), 317–347.

[20] W. Van Assche, *Nearest neighbor recurrence relations for multiple orthogonal polynomials*, J. Approx. Theory **163** (2011), no. 10, 1427–1448.

[21] W. Van Assche, *Ratio asymptotics for multiple orthogonal polynomials*, Contemporary Mathematics (to appear); arXiv:1408.1829 [math.CA].

Walter Van Assche
Department of Mathematics
KU Leuven
Celestijnenlaan 200 B box 2400
BE-3001 Leuven
Belgium
`walter@wis.kuleuven.be`

Thorsten Neuschel
current address:
IRMP
Université Catholique de Louvain
Chemin du Cyclotron 2
BE-1348 Louvain-la-Neuve
Belgium
`thorsten.neuschel@uclouvain.be`